



INDEX PRONOSTIQUES

Deux index sont proposés dans la littérature :

1. L'index DS-GPA (*disease specific-graded prognostic assessment*)

Il a été publié en 2012 par Sperduto *et al.* (4). Score établi à partir d'une série de 3666 CBNPC et 562 CPC, données recueillies entre 1985 et 2007. Il utilise le score GPA détaillé dans le tableau ci-dessous, avec des survies analysées spécifiquement dans les cancers bronchiques : index DS-GPA ou « Lung GPA »

	SCORE GPA		
	0	0,5	1
Age	> 60 ans	50-59 ans	< 50 ans
KPS	< 70	70-80	90-100
Nombre de métastase cérébrale	> 3	2 à 3	1
Métastases extra crâniennes	Oui	–	Non

Valeur du Score DS-GPA	Survie médiane CBNPC (mois)	Survie médiane CPC (mois)
3,5-4	14,78	2,79
2,5-3	9,43	4,5
1,5-2	5,49	4,9
0-1	3,02	2,7
Tous	7	4,9

Tableau 1- Index DS Lung GPA

Le tableau 1 montre les médianes de survie spécifiques pour les métastases cérébrales des cancers bronchiques : elles varient de 2,7 à 14,8 mois. Les facteurs pronostiques retrouvés sont l'âge, l'index de Karnofsky, le nombre de métastase cérébrale et la présence ou non de métastase extra cérébrale. Ce score est le plus communément admis pour évaluer le pronostic des patients avec des métastases cérébrales d'un cancer bronchique.



2. L'index Lung-molGPA (*Lung-molecular-graded prognostic assessment*)

Ce nouveau score publié en 2017 (5) élaboré à partir de 2186 patients, concerne uniquement les CBNPC, en les séparant en 2 groupes : adénocarcinomes et non adénocarcinomes. Par ailleurs ce score analyse des données plus récentes, recueillies entre 2006 et 2014. De plus, il ajoute dans les facteurs pronostiques la présence d'une mutation de l'*EGFR* ou d'un réarrangement d'*ALK*, qui comptent comme 1 point dans le score. Ainsi, seuls les patients avec une mutation de l'*EGFR* ou un réarrangement d'*ALK* peuvent avoir un score à 4 (Tableau 2).

Score	0	0,5	1
Age	≥70 ans	< 70 ans	////
KPS	< 70	80	90-100
Métastases extra crâniennes	OUI	////	NON
Nombre de MC	>4	1 à 4	////
Statut oncogénique	Pas d'altération <i>EGFR</i> ou <i>ALK</i>	////	Altération <i>EGFR</i> ou <i>ALK</i>

Valeur du score Lung-molGPA	Type histologique	0-1	1,5-2,5	2,5-3,5	3,5-4	Tout
Médiane de survie globale (mois)	Non ADK	5,3	9,8	12,8	NA	9,2
	ADK	6,9	13,7	26,5	46,8	15,2

Tableau 2- Score Lung-molGPA : calcul (haut) et survie globale selon la valeur du score.

Les survies globales selon ce score montrent des survies significativement plus élevées que le score DS-GPA, atteignant 46 mois pour les patients avec un score entre 3,5 à 4 et un ADK (ce groupe comprend les patients avec mutation *EGFR* ou réarrangement de *ALK*). Il faut donc utiliser ce score de préférence pour les patients avec un CBNPC, surtout chez les patients avec une addiction oncogénique *EGFR* ou *ALK*.



REFERENCES

1. Alexander BM, Brown PD, Ahluwalia MS, Aoyama H, Baumert BG, Chang SM, et al. Clinical trial design for local therapies for brain metastases: a guideline by the Response Assessment in Neuro-Oncology Brain Metastases working group. *Lancet Oncol.* 2018;19(1):e33–e42.
2. Peters S, Camidge DR, Shaw AT, Gadgeel S, Ahn JS, Kim D-W, et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med.* 31 2017;377(9):829-38.
3. Chargari C, Dhermain F. Métastases cérébrales d'un cancer bronchique non à petites cellules: du traitement standardisé au traitement personnalisé. *Rev Mal Respir Actual.* 2013;5(5):547–556.
4. Sperduto PW, Kased N, Roberge D, Xu Z, Shanley R, Luo X, et al. Summary report on the graded prognostic assessment: an accurate and facile diagnosis-specific tool to estimate survival for patients with brain metastases. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 févr 2012;30(4):419-25.
5. Sperduto PW, Yang TJ, Beal K, Pan H, Brown PD, Bangdiwala A, et al. Estimating Survival in Patients With Lung Cancer and Brain Metastases: An Update of the Graded Prognostic Assessment for Lung Cancer Using Molecular Markers (Lung-molGPA). *JAMA Oncol.* 1 juin 2017;3(6):827-31.
6. Schellinger PD, Meinck HM, Thron A. Diagnostic accuracy of MRI compared to CCT in patients with brain metastases. *J Neurooncol.* 1999;44(3):275-81.
7. Sighvatsson V, Ericson K, Tómasson H. Optimising contrast-enhanced cranial CT for detection of brain metastases. *Acta Radiol Stockh Swed* 1987. nov 1998;39(6):718-22.
8. Seute T, Leffers P, ten Velde GPM, Twijnstra A. Detection of brain metastases from small cell lung cancer: consequences of changing imaging techniques (CT versus MRI). *Cancer.* 15 avr 2008;112(8):1827-34.
9. Olson JJ, Kalkanis SN, Ryken TC. Congress of Neurological Surgeons Systematic Review and Evidence-Based Guidelines for the Treatment of Adults With Metastatic Brain Tumors: Executive Summary. *Neurosurgery.* 01 2019;84(3):550-2.
10. Patchell RA, Tibbs PA, Walsh JW, Dempsey RJ, Maruyama Y, Kryscio RJ, et al. A randomized trial of surgery in the treatment of single metastases to the brain. *N Engl J Med.* 22 févr 1990;322(8):494-500.
11. Noordijk EM, Vecht CJ, Haaxma-Reiche H, Padberg GW, Voormolen JH, Hoekstra FH, et al. The choice of treatment of single brain metastasis should be based on extracranial tumor activity and age. *Int J Radiat Oncol Biol Phys.* 1 juill 1994;29(4):711-7.
12. Mintz AH, Kestle J, Rathbone MP, Gaspar L, Hugenholtz H, Fisher B, et al. A randomized trial to assess the efficacy of surgery in addition to radiotherapy in patients with a single cerebral metastasis. *Cancer.* 1 oct 1996;78(7):1470-6.
13. Latorzeff I, Antoni D, Gaudaire-Josset S, Feuvret L, Tallet-Richard A, Truc G, et al. Radiothérapie des métastases cérébrales. *Cancer/Radiothérapie.* sept 2016;20:S80-7.
14. Sahgal A, Aoyama H, Kocher M, Neupane B, Collette S, Tago M, et al. Phase 3 trials of stereotactic radiosurgery with or without whole-brain radiation therapy for 1 to 4 brain metastases: individual patient data meta-analysis. *Int J Radiat Oncol Biol Phys.* 15 mars 2015;91(4):710-7.
15. Gerosa M, Nicolato A, Foroni R, Tomazzoli L, Bricolo A. Analysis of long-term outcomes and prognostic factors in patients with non-small cell lung cancer brain metastases treated by gamma knife radiosurgery. *J Neurosurg.* janv 2005;102 Suppl:75-80.
16. Motta M, del Vecchio A, Attuati L, Picozzi P, Perna L, Franzin A, et al. Gamma knife radiosurgery for treatment of cerebral metastases from non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys.* 15 nov 2011;81(4):e463-468.
17. Fuentes R, Osorio D, Expósito Hernandez J, Simancas-Racines D, Martínez-Zapata MJ, Bonfill Cosp X. Surgery versus stereotactic radiotherapy for people with single or solitary brain metastasis. *Cochrane Database Syst Rev.* 20 2018;8:CD012086.



18. Yamamoto M, Serizawa T, Shuto T, Akabane A, Higuchi Y, Kawagishi J, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLGK0901): a multi-institutional prospective observational study. *Lancet Oncol.* avr 2014;15(4):387-95.
19. Shuto T, Akabane A, Yamamoto M, Serizawa T, Higuchi Y, Sato Y, et al. Multiinstitutional prospective observational study of stereotactic radiosurgery for patients with multiple brain metastases from non-small cell lung cancer (JLGK0901 study-NSCLC). *J Neurosurg.* 01 2018;129(Suppl1):86-94.
20. Le Rhun É, Dhermain F, Noël G, Reyns N, Carpentier A, Mandonnet E, et al. [ANOCEF guidelines for the management of brain metastases]. *Cancer Radiother J Soc Francaise Radiother Oncol.* févr 2015;19(1):66-71.
21. Brown PD, Jaeckle K, Ballman KV, Farace E, Cerhan JH, Anderson SK, et al. Effect of Radiosurgery Alone vs Radiosurgery With Whole Brain Radiation Therapy on Cognitive Function in Patients With 1 to 3 Brain Metastases: A Randomized Clinical Trial. *JAMA.* 26 juill 2016;316(4):401-9.
22. Mahajan A, Ahmed S, McAleer MF, Weinberg JS, Li J, Brown P, et al. Post-operative stereotactic radiosurgery versus observation for completely resected brain metastases: a single-centre, randomised, controlled, phase 3 trial. *Lancet Oncol.* août 2017;18(8):1040-8.
23. Lamba N, Muskens IS, DiRisio AC, Meijer L, Briceno V, Edrees H, et al. Stereotactic radiosurgery versus whole-brain radiotherapy after intracranial metastasis resection: a systematic review and meta-analysis. *Radiat Oncol Lond Engl.* 24 juin 2017;12(1):106.
24. Soffietti R, Kocher M, Abacioglu UM, Villa S, Fauchon F, Baumert BG, et al. A European Organisation for Research and Treatment of Cancer phase III trial of adjuvant whole-brain radiotherapy versus observation in patients with one to three brain metastases from solid tumors after surgical resection or radiosurgery: quality-of-life results. *J Clin Oncol Off J Am Soc Clin Oncol.* 1 janv 2013;31(1):65-72.
25. Brown PD, Ballman KV, Cerhan JH, Anderson SK, Carrero XW, Whitton AC, et al. Postoperative stereotactic radiosurgery compared with whole brain radiotherapy for resected metastatic brain disease (NCCTG N107C/CEC-3): a multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol.* 2017;18(8):1049-60.
26. Ma TM, Grimm J, McIntyre R, Anderson-Keightly H, Kleinberg LR, Hales RK, et al. A prospective evaluation of hippocampal radiation dose volume effects and memory deficits following cranial irradiation. *Radiother Oncol J Eur Soc Ther Radiol Oncol.* 2017;125(2):234-40.
27. Rodríguez de Dios N, Couñago F, López JL, Calvo P, Murcia M, Rico M, et al. Treatment Design and Rationale for a Randomized Trial of Prophylactic Cranial Irradiation With or Without Hippocampal Avoidance for SCLC: PREMIER Trial on Behalf of the Oncologic Group for the Study of Lung Cancer/Spanish Radiation Oncology Group-Radiation Oncology Clinical Research Group. *Clin Lung Cancer.* sept 2018;19(5):e693-7.
28. Zindler JD, Bruynzeel AME, Eekers DBP, Hurkmans CW, Swinnen A, Lambin P. Whole brain radiotherapy versus stereotactic radiosurgery for 4-10 brain metastases: a phase III randomised multicentre trial. *BMC Cancer.* 25 juill 2017;17(1):500.
29. Roberge D, Brown PD, Whitton A, O'Callaghan C, Leis A, Greenspoon J, et al. The Future Is Now-Prospective Study of Radiosurgery for More Than 4 Brain Metastases to Start in 2018! *Front Oncol.* 2018;8:380.
30. Robinet G, Thomas P, Breton JL, Léna H, Gouva S, Dabouis G, et al. Results of a phase III study of early versus delayed whole brain radiotherapy with concurrent cisplatin and vinorelbine combination in inoperable brain metastasis of non-small-cell lung cancer: Groupe Français de Pneumo-Cancérologie (GFPC) Protocol 95-1. *Ann Oncol Off J Eur Soc Med Oncol ESMO.* janv 2001;12(1):59-67.
31. Barlesi F, Gervais R, Lena H, Hureaux J, Berard H, Paillot D, et al. Pemetrexed and cisplatin as first-line chemotherapy for advanced non-small-cell lung cancer (NSCLC) with asymptomatic inoperable brain metastases: a multicenter phase II trial (GFPC 07-01). *Ann Oncol Off J Eur Soc Med Oncol ESMO.* nov 2011;22(11):2466-70.
32. Besse B, Lasserre SF, Compton P, Huang J, Augustus S, Rohr U-P. Bevacizumab safety in patients with central nervous system metastases. *Clin Cancer Res Off J Am Assoc Cancer Res.* 1 janv 2010;16(1):269-78.



33. Socinski MA, Langer CJ, Huang JE, Kolb MM, Compton P, Wang L, et al. Safety of bevacizumab in patients with non-small-cell lung cancer and brain metastases. *J Clin Oncol Off J Am Soc Clin Oncol*. 1 nov 2009;27(31):5255-61.
34. Besse B, Le Moulec S, Mazières J, Senellart H, Barlesi F, Chouaid C, et al. Bevacizumab in Patients with Nonsquamous Non-Small Cell Lung Cancer and Asymptomatic, Untreated Brain Metastases (BRAIN): A Nonrandomized, Phase II Study. *Clin Cancer Res Off J Am Assoc Cancer Res*. 15 avr 2015;21(8):1896-903.
35. Brahmer J, Reckamp KL, Baas P, Crinò L, Eberhardt WEE, Poddubskaya E, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *N Engl J Med*. 9 juill 2015;373(2):123-35.
36. Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. *N Engl J Med*. 27 sept 2015;
37. Goldman JW, Crino L, Vokes EE, Holgado E, Reckamp K, Pluzanski A, et al. P2.36: Nivolumab (nivo) in Patients (pts) With Advanced (adv) NSCLC and Central Nervous System (CNS) Metastases (mets): Track: Immunotherapy. *J Thorac Oncol Off Publ Int Assoc Study Lung Cancer*. oct 2016;11(10S):S238-9.
38. Hellmann MD, Paz-Ares L, Bernabe Caro R, Zurawski B, Kim S-W, Carcereny Costa E, et al. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 28 sept 2019;
39. Goldberg SB, Gettinger SN, Mahajan A, Chiang AC, Herbst RS, Sznol M, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol*. juill 2016;17(7):976-83.
40. Gandhi L, Rodríguez-Abreu D, Gadgeel S, Esteban E, Felip E, De Angelis F, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. *N Engl J Med*. 31 mai 2018;378(22):2078-92.
41. Rittmeyer A, Barlesi F, Waterkamp D, Park K, Ciardiello F, von Pawel J, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet Lond Engl*. 12 déc 2016;
42. Hendriks LEL, Henon C, Auclin E, Mezquita L, Ferrara R, Audigier-Valette C, et al. Outcome of Patients with Non-Small Cell Lung Cancer and Brain Metastases Treated with Checkpoint Inhibitors. *J Thorac Oncol*. juill 2019;14(7):1244-54.
43. Tallet AV, Dhermain F, Le Rhun E, Noël G, Kirova YM. Combined irradiation and targeted therapy or immune checkpoint blockade in brain metastases: toxicities and efficacy. *Ann Oncol Off J Eur Soc Med Oncol*. 1 déc 2017;28(12):2962-76.
44. Shepard MJ, Xu Z, Donahue J, Eluvathingal Muttikkal TJ, Cordeiro D, Hansen L, et al. Stereotactic radiosurgery with and without checkpoint inhibition for patients with metastatic non-small cell lung cancer to the brain: a matched cohort study. *J Neurosurg*. 26 juill 2019;1-8.
45. Chen L, Douglass J, Kleinberg L, Ye X, Marciscano AE, Forde PM, et al. Concurrent Immune Checkpoint Inhibitors and Stereotactic Radiosurgery for Brain Metastases in Non-Small Cell Lung Cancer, Melanoma, and Renal Cell Carcinoma. *Int J Radiat Oncol Biol Phys*. 15 mars 2018;100(4):916-25.
46. Linskey ME, Andrews DW, Asher AL, Burri SH, Kondziolka D, Robinson PD, et al. The role of stereotactic radiosurgery in the management of patients with newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. *J Neurooncol*. janv 2010;96(1):45-68.
47. Langley RE, Stephens RJ, Nankivell M, Pugh C, Moore B, Navani N, et al. Interim data from the Medical Research Council QUARTZ Trial: does whole brain radiotherapy affect the survival and quality of life of patients with brain metastases from non-small cell lung cancer? *Clin Oncol R Coll Radiol G B*. mars 2013;25(3):e23-30.
48. Mulvenna P, Nankivell M, Barton R, Faivre-Finn C, Wilson P, McColl E, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet Lond Engl*. 22 oct 2016;388(10055):2004-14.
49. Soria J-C, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med*. 18 nov 2017;



50. Ramalingam SS, Vansteenkiste J, Planchard D, Cho BC, Gray JE, Ohe Y, et al. Overall Survival with Osimertinib in Untreated, EGFR-Mutated Advanced NSCLC. *N Engl J Med*. 02 2020;382(1):41-50.
51. Magnuson WJ, Lester-Coll NH, Wu AJ, Yang TJ, Lockney NA, Gerber NK, et al. Management of Brain Metastases in Tyrosine Kinase Inhibitor-Naïve Epidermal Growth Factor Receptor-Mutant Non-Small-Cell Lung Cancer: A Retrospective Multi-Institutional Analysis. *J Clin Oncol Off J Am Soc Clin Oncol*. 1 avr 2017;35(10):1070-7.
52. Miyawaki E, Kenmotsu H, Mori K, Harada H, Mitsuya K, Mamesaya N, et al. Optimal Sequence of Local and EGFR-TKI Therapy for EGFR-Mutant Non-Small Cell Lung Cancer With Brain Metastases Stratified by Number of Brain Metastases. *Int J Radiat Oncol Biol Phys*. 01 2019;104(3):604-13.
53. Lee C-C, Hsu SPC, Lin C-J, Wu H-M, Chen Y-W, Luo Y-H, et al. Epidermal growth factor receptor mutations: association with favorable local tumor control following Gamma Knife radiosurgery in patients with non-small cell lung cancer and brain metastases. *J Neurosurg*. 21 juin 2019;1-8.
54. Reungwetwattana T, Nakagawa K, Cho BC, Cobo M, Cho EK, Bertolini A, et al. CNS Response to Osimertinib Versus Standard Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Patients With Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol Off J Am Soc Clin Oncol*. 28 août 2018;JCO2018783118.
55. Ballard P, Yates JWT, Yang Z, Kim D-W, Yang JC-H, Cantarini M, et al. Preclinical Comparison of Osimertinib with Other EGFR-TKIs in EGFR-Mutant NSCLC Brain Metastases Models, and Early Evidence of Clinical Brain Metastases Activity. *Clin Cancer Res Off J Am Assoc Cancer Res*. 15 oct 2016;22(20):5130-40.
56. Mok TS, Wu Y-L, Ahn M-J, Garassino MC, Kim HR, Ramalingam SS, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med*. 6 déc 2016;
57. Wang X, Xu Y, Tang W, Liu L. Efficacy and Safety of Radiotherapy Plus EGFR-TKIs in NSCLC Patients with Brain Metastases: A Meta-Analysis of Published Data. *Transl Oncol*. oct 2018;11(5):1119-27.
58. Chang EL, Wefel JS, Hess KR, Allen PK, Lang FF, Kornguth DG, et al. Neurocognition in patients with brain metastases treated with radiosurgery or radiosurgery plus whole-brain irradiation: a randomised controlled trial. *Lancet Oncol*. nov 2009;10(11):1037-44.
59. Soffietti R, Kocher M, Abacioglu UM, Villa S, Fauchon F, Baumert BG, et al. A European Organisation for Research and Treatment of Cancer phase III trial of adjuvant whole-brain radiotherapy versus observation in patients with one to three brain metastases from solid tumors after surgical resection or radiosurgery: quality-of-life results. *J Clin Oncol Off J Am Soc Clin Oncol*. 1 janv 2013;31(1):65-72.
60. Soria J-C, Tan DSW, Chiari R, Wu Y-L, Paz-Ares L, Wolf J, et al. First-line ceritinib versus platinum-based chemotherapy in advanced ALK-rearranged non-small-cell lung cancer (ASCEND-4): a randomised, open-label, phase 3 study. *Lancet Lond Engl*. 4 mars 2017;389(10072):917-29.
61. Solomon BJ, Mok T, Kim D-W, Wu Y-L, Nakagawa K, Mekhail T, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med*. 4 déc 2014;371(23):2167-77.
62. Camidge DR, Kim D-W, Tiseo M, Langer CJ, Ahn M-J, Shaw AT, et al. Exploratory Analysis of Brigatinib Activity in Patients With Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer and Brain Metastases in Two Clinical Trials. *J Clin Oncol Off J Am Soc Clin Oncol*. 10 2018;36(26):2693-701.
63. Horn L, Mansfield AS, Szczesna A, Havel L, Krzakowski M, Hochmair MJ, et al. First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer. *N Engl J Med*. 6 déc 2018;379(23):2220-9.